



## A survey of the bushmeat trade of the straw-coloured fruit bat (*Eidolon helvum* Kerr, 1792) at Maele Island (Kisangani city, Democratic Republic of the Congo)

Musaba P. Akawa<sup>1</sup>, Malekani A. Bendeki<sup>2</sup>, Kirongozi F. Botelanyele<sup>2</sup>, Shabani I<sup>1</sup>, Nebesse C. Mololo<sup>1</sup>, Van V. Cakenberghe<sup>3</sup>, E. Verheyen<sup>4,5</sup>; Gembu G.C. Tungaluna<sup>1,2</sup>, Justin A. Asimonyio<sup>2</sup>, Masengo C. Ashande<sup>6</sup>, Koto-te-Nyiwa Ngbolua<sup>6,7,8</sup>

<sup>1</sup>Université de Kisangani, Faculté des Sciences, Département d'Ecologie et de Gestion de Ressources Animales, B.P. 2012 Kisangani, République Démocratique du Congo

<sup>2</sup>Université de Kisangani, Centre de Surveillance de la Biodiversité, B.P. 2012 Kisangani, République Démocratique du Congo

<sup>3</sup>University of Antwerp, Functional Morphology Laboratory, Antwerpen, Belgium

<sup>4</sup>University of Antwerp, Evolutionary Ecology Group, Antwerpen, Belgium

<sup>5</sup>Royal Belgian Institute of Natural Sciences, OD Taxonomy and Phylogeny, Brussels, Belgium

<sup>6</sup>Université de Gbado-Lite, Faculté des Sciences, Département des Sciences de l'Environnement, B.P. 111 Gbado-Lite, Province de Nord Ubangi, République Démocratique du Congo

<sup>7</sup>Institut Supérieur Pédagogique d'Abumombazi, Abumombazi, Province de Nord Ubangi, République Démocratique du Congo

<sup>8</sup>Université de Kinshasa, Faculté des Sciences, Département de Biologie, B.P. 190 Kinshasa XI, République Démocratique du Congo

\*Corresponding author: Koto-te-Nyiwa Ngbolua, E-mail: [jpngbolua@unikin.ac.cd](mailto:jpngbolua@unikin.ac.cd)

Received: March 24, 2017, Accepted: April 15, 2017, Published: April 15, 2017.

### ABSTRACT:

Although bats carry out important beneficial ecological and agricultural functions such as pollination and dispersion of seeds, they often carry dangerous diseases. This is worrying because of the increased dependency of the urban populations in Kisangani on bats as a source of animal proteins. The straw-coloured fruit bat (*Eidolon helvum*) is one of the most notable bat species that is both important as a food source and as a host for zoonotic diseases. It is a widely distributed species, which is relatively common throughout its distribution area. Recently, it was classified as Near Threatened on the IUCN Red List due to a decreasing population trend. To evaluate the hunting pressure on this species and the supply of straw-coloured fruit bats for human consumption, we inventoried the carcasses of this species on one of the main bushmeat markets in the region of Kisangani, situated on the island of Maele. Our survey was carried out between January and December 2013 and targeted hunters as well as traders. In total, we counted 3,034 carcasses. Our preliminary results suggest that the hunting pressure on the straw-coloured fruit bat is rather stable. It is highest during the months of October, September and May and lowest in April and December. Our results do not show a decline in trade of these bats during August through November, when hunting is illegal in the DR Congo. This suggests that hunting activities are not ceased during that period. The fact that the local population consumes these bats on a regular basis throughout the year creates a very important opportunity for spill-over events that may lead to the outbreaks of zoonotic diseases.

**Keyword:** Bat, bushmeat, hunting, *Eidolon helvum*, Democratic Republic of the Congo

### INTRODUCTION

The Democratic Republic of the Congo is a biogeographic area of high value for the conservation of the biodiversity and is very rich in plant and animal species [1-16]. Among the terrestrial vertebrates, the bats are one of the most abundant and diversified ecological groups. They belong to the mammalian order Chiroptera, which includes a large variety of species, second only to the Rodents. Worldwide, there are approximately 1,300 bat species representing about one quarter of all known mammals [17].

Bats are part of the wildlife, which is very sought for as bushmeat by societies living along the rivers in forests, such as the area around Kisangani. As such, they contribute a major part to the animal protein consumed by local communities and therefore become a source of income for numerous families of resident hunters [18], who hunt these bats year-round, neglecting the closed season.

The fierce exploitation of the frugivorous *Eidolon helvum* has a major impact on an ecological level, given the fact that this species

fulfills a number of roles, including the dispersion of seeds, needed to regenerate the forests in various countries in Africa and the Paleotropics [19]. Some of these seeds are of plants with an important economic value, e.g. *Micilia excelsa* [20]. The excessive killing of *E. helvum* can endanger its capacity to provide these vital ecosystem services [21].

In Kisangani, bats are hunted for food, for not only personal use, but also for commercial reasons. Information on quantities or yields is completely lacking and hardly anything is known about the proportion of fruit-bat carcasses.

### MATERIAL AND METHODS

#### Study area


Individual specimens belonging to *Eidolon helvum* (Kerr, 1792) were inventoried on the island of Maele, within the city of Kisangani, Tshopo Province, Democratic Republic of the Congo (DRC) (00° 29' 19.3" N 25° 12' 42.8" E, 376 m). It is located downstream of the island of Mbiye and upstream of the Wagenya Falls fishery, a tourist attraction near the city of Kisangani.

Situated in the equatorial zone, Maele island globally has the same climate as the city of Kisangani: Af (Köppen classification type: Tropical Rain forest [22]. The mean temperature of the coldest month is above 18 °C, and the annual temperature variation is less than 5 °C. The average rainfall during the driest month is about 60 mm [23]. In Kisangani, the rains are abundant over the year with an unequal monthly distribution, which results in two very wet seasons ("wet": from March to May and from September to November) and two seasons with lower rain intensities ("dry": December to February and June to August) [24].

Maele Island has a mainly man-made landscape. Its original vegetation is almost completely lost. From the north to the south and from the east to the west it is dominated by fields of various cultures. There is no primary nor secondary forest to speak of, only a few small young secondary forest islands dominated by *Myrianthus arboreus* P. Beauv., 1805 (Urticaceae - Giant yellow mulberry), *Pseudospondiace microcarpa* A. Rich., 1883 (Anacardiaceae - African Grape) *Musanga cecropioides* R. Brown ex Tedlie (Urticaceae - African corkwood tree or umbrella tree) and a few budding *Piptadeniastrum africanum* (Hook.f.) Brenan, 1955 (Fabaceae - African greenheart).

The fallows are dominated by lianes: *Chromolaena odorata* (L.) King & H.E. Robins (Asteraceae - Siam weed, Christmas bush, devil weed, camfhur grass, common floss flower and trifid), *Triumfetta cordifolia* A. Rich (Malvaceae - cordleaf burrbark), and *Alcornea floribunda* Müll. Arg. (Euphorbiaceae).

Furthermore, there are numerous fields with various crops: rice, corn, bananas, vegetables, cassava, etc.



**Figure 1:** Map showing the study area with the location of the Maele Island (IMA) where the bats were hunted and opposite Maele market (MMA), where the bats were sold.

### Hunting Method

Hunting on Maele Island is done during the entire year and the hunters use mist nets to capture bats belonging to various species. We were interested in *Eidolon helvum*, which is not only the most hunted species, but also the most abundant one and the one, which is most on sale on the market in Kisangani.

The hunters use canoes, which take them to the island in about 10 minutes.



**Figure 2:** (a) A bat hunter returning in his canoe from the island

of Maele near the city of Kisangani. (b) A female trader waiting for the return of the hunter.

### Inventory of the carcasses


3,034 carcasses of *Eidolon helvum* captured on Maele Island were inventoried on the market at Maele between January and December 2013. During that period, we were present on the landing site before the arrival of the first hunter and remained there until after the arrival of the last hunter.

At each arrival of a hunter, we checked the specimens belonging to *Eidolon helvum* and noted the following parameters: number of specimens, sex, age, weight and price. Afterwards, we also counted the number of bat specimens that were available at the trader's displays to obtain an estimate of the daily supply.

Every morning, between 7 and 10 a.m., the traders buy the bats from the hunters and they sell them in the evening. All carcasses are boiled before being sold and bats, that were not sold, are stocked in a freezer to be sold the next day.


### RESULTS

The figure 3 gives an overview on the monthly number of *Eidolon helvum* bats on sale at the Maele Island market based on the daily data obtained from the hunters and traders.



**Figure 3:** Overview of the monthly number of *Eidolon helvum* bats on sale at the Maele Island market year-round.

The figure shows that there was an extremely large amount of bats available in October (710 specimens), which was building up from August through September (386). This peak coincides with one of the wet seasons. A second, much smaller peak overlaps with end of the second wet season in May (349 bats). Remarkably, however, the number of bats available on the market in both March and April was low.




**Figure 4:** Estimated number of sold bats on the Maele island market, divided by sex and age.

From the figure 4, when the monthly numbers are split up by sex and age, it is clear that an overwhelming number of sold bats were


non-adults, especially during the peak of the wet season (from September to November). Adult bats were only available in relatively high proportions in January and February. Throughout the year, females were generally more available than males, only in May (and to a lesser extent in March) there were more males than females.

The figure 5 gives the distribution of the availability of straw-coloured fruit bats on the market at Maele Island over the different seasons.



**Figure 5:** Frequencies according to the seasons (DJF: December - January -February; MAM: March-April-May; JJA: June-July-August; SON: September-October-November).

The graph shows a steady built-up of the number of bats available on the market from the December to February dry season, over the wet season from March to May and the second dry season from June to August. Most bats were available during the second wet season of the year: from September to November, when 1,318 carcasses were inventoried (43.63 % of all specimens).



**Figure 6:** Ratio of the bats on sale during the period that hunting is allowed (PBHI: December through July) or not (PDHI: August through November).

In the Kisangani area, hunting bats is allowed from December through July (PBHI: Period before the hunting interdiction). However, most bat carcasses (53%) were sold during the period that the hunt and the trade in bushmeat was illegal (PDHI: Period during the hunting interdiction) (Figure 6).

## DISCUSSION

During survey period, 3,034 specimens or carcasses of *Eidolon helvum* were inventoried on the market at Maele, where

they were sold as bushmeat. The abundance of *E. helvum* among the bats available on the market at Maele is also linked to the species' habits: roosting in trees by the thousands. The genus *Eidolon* is known for the large groups it forms, consisting of thousands or millions of individuals [25-27]. As such, they offer an opportunity for hunters to kill them in large numbers. Furthermore, being of small size, they are easily transportable to the markets. This explains why these bats are found in large numbers on markets.

Additionally, the lack of alternative steady income-generating activities for the majority of the hunters and the consumer's taste for these bats generate an extra pressure to capture *E. helvum* bats. During the current study, besides the 3,034 *E. helvum* specimens, 147 *Hypsignathus monstrosus* H. Allen 1861, 51 *Myonycteris torquata* (Dobson, 1878), 19 *Rousettus aegyptiacus* (E. Geoffroy, 1810), and 15 *Epomops franqueti* (Tomes, 1860) were registered. A study by [28] on the seasonal distribution of fruit bats at Yoko in relation to the various seasons showed that the capture of Pteropodids was only profitable during the season with abundant rains. During this season, 59% of the *Casinycteris argynnis* Thomas, 1910 specimens were captured, as well as 79% of *Epomops franqueti*, 95% of *Hypsignathus monstrosus*, 91% of *Myonycteris torquata* and 69% of all *Scotonycteris zenkeri* Matschie, 1894 [= *Scotonycteris bergmansi* [29]]. In the seasons with lower rain intensities, the only species that were more or less profitable to collect were *Megaloglossus woermanni* Pagenstecher, 1885 (71% of all captures) and *Rousettus aegyptiacus* (58% of all captures).

With the exception of [30], all studies covering of bats on the main market at Kisangani mention *Eidolon helvum* to be present. The author [31] attributed the high rank of *E. helvum* in the local food supply to its large size. This makes it a highly valued species for the hunters. Furthermore, [32] found that 43.2% of all the fresh mammals sold on the market in 1989 were representatives of this bat species: 2,005 out of 4,637 (the second most numerous mammalian order were the Primates).

Bushmeat is an important source of animal protein in the daily diet of populations in the Congo Basin [33], and its contribution to the household economy is significant [18, 34].

Although hunting in the DRC is covered by law (laws 82-002 of 28 May 1982, 014/CAB/MIN/ENV/2004 of 29 April 2004 and 14/003 of 11 February 2014 [35] which closes the hunt during certain periods of the year, it is often condoned by local authorities as is illustrated in Figure 6. The author [36] claimed that the closed hunting season was abided until the early 1970s, after which it watered down. The intense poaching on *E. helvum*, however, might lead to a faster extinction in view of the low reproduction rate of this species [36-38]. In Kisangani, economic profit unfortunately does not care at all about the sustainability of the species, hence the game market is operational year-round, and wildlife is subject to continuous hunting pressure throughout the year.

Jenkins *et al.* [39], who discussed the consumption of bat meat in Madagascar, demonstrated the appreciation of this type of meat by the local population. The preferred species included three fruit bats (*Pteropus rufus* Tiedemann, 1808 [= *P. niger* Kerr, 1792], *Eidolon dupreanum* (Pollen, 1867), and *Rousettus madagascariensis* G. Grandidier, 1929) and one large-sized insect-eating bat: *Hipposideros commersoni* (E. Geoffroy, 1813). Additionally, they also reported the consumption of small insect-eating bats. As in Kisangani, they also found that the hunting season is generally not respected, as bats were also available during the months that hunting was illegal. In some areas of

Madagascar, this causes a serious threat on bat populations. They also found little data on the socioeconomic or cultural values of bats.

In Yoko, the author [28] found that the majority of specimens captured were adults or subadults, the present study found that an overwhelming part of the specimens presented for sale at the Maele market were non-adult specimens. Unfortunately, we do not have any detailed information concerning their exact age category (juvenile, immature, sub-adult), so they might represent various stages in their development. Nevertheless, given the fact that *E. helvum* only has one young a year and that the number of males and females are more or less equal [27], it remains surprising to see that such a large amount of non-adults were captured. This might possibly be the result of these young (er) bats being less agile or less able to avoid the nets being used to capture them. Or perhaps they might be captured purposely because of the local food taste. The author [27] also mention "At Kampala, rain falls throughout the year but there are peaks in Oct-Nov and Apr-May, and the reproductive chronology is such that both spermatogenesis and lactation coincide with periods of high rainfall, and births during the relatively dry periods are avoided." This is very comparable with the situation in Kisangani and would explain the high peak in the September through November wet season.

## CONCLUSION

During the year 2013, 3,034 *Eidolon helvum* carcasses were inventoried on the market at Maele, right across Maele Island, where these animals were captured.

The hunting pressure on Maele Island was highest during the months of October, September and May with respectively 710, 386 and 349 carcasses. Females formed the majority of bats captured during almost the entire year, especially during the months of September and October (wet season). Males were only in the majority in May. Non-adult bats were dominantly present from May to December, but also in lower numbers in March. The only month that adults were in the majority was January, although there were still a large number of non-adult bats available.

Although there is a major fluctuation, hunting occurs during the entire year, implying that the closure of the hunting season from August through November is not respected in the area around Kisangani especially as most of the carcasses were on sale during that period.

## ACKNOWLEDGEMENTS

The authors are indebted to the people who participated in this work: Mr. Moise Bipoo, Mr. Marc Boketshu, Mr. Bienvenu Ndjoku, and to the hunters and traders for their important contribution in the realization of this study.

## REFERENCES

1. J.A. Asimonyio, K. Kambale, E. Shutsha, G.N. Bongo, D.S.T. Tshibangu, P.T. Mpiana, K.N. Ngbolua. Phytoecological Study of Uma Forest (Kisangani City, Democratic Republic Of The Congo). *J. of Advanced Botany and Zoology*, V312 (2015). DOI: 10.15297/JABZ.V312.01.
2. J.A. Asimonyio, J.C. Ngabu, C.B. Lomba, C.M. Falanga, P.T. Mpiana, K.N. Ngbolua. Structure et diversité d'un peuplement forestier hétérogène dans le bloc sud de la réserve forestière de Yoko (Ubundu, République Démocratique du Congo). *International Journal of Innovation and Scientific Research* 18 (2) (2015) 241-251.
3. B.G. Badjedjea, B.J. Akuboy, M.F. Masudi, J.A. Asimonyio, K.P. Museu, K.N. Ngbolua. A preliminary survey of the amphibian fauna of Kisangani eco-region, Democratic Republic of the Congo. *J. of Advanced Botany and Zoology*, V314 (2015). DOI: 10.15297/JABZ.V314.01.
4. P. Baelo, J.A. Asimonyio, S. Gambalemoke, N. Amundala, R. Kiakenya, E. Verheyen, A. Laudisoit, K.N. Ngbolua. Reproduction et structure des populations des Sciuridae (Rodentia, Mammalia) de la réserve forestière de Yoko (Ubundu, RD Congo). *International Journal of Innovation and Scientific Research* 23 (2) (2016) 428-442.
5. P. Baelo, C. Kahandi, J. Akuboyi1, J.L. Juakaly, K.N. Ngbolua. Contribution à l'étude de la biodiversité et de l'écologie des Araignées du sol dans un champ cultivé de *Manihot esculenta* Crantz (Euphorbiaceae) à Kisangani, RD Congo. *International Journal of Innovation and Scientific Research* 23 (2) (2016) 412-418.
6. J.K. Kambale, F.M. Feza, J.M. Tsongo, J.A. Asimonyio, S. Mapeta, H. Nshimba, B.Z. Gbolo, P.T. Mpiana, K.N. Ngbolua. La filière bois-énergie et dégradation des écosystèmes forestiers en milieu périurbain: Enjeux et incidence sur les riverains de l'île Mbiye à Kisangani (République Démocratique du Congo). *International Journal of Innovation and Scientific Research* 21 (1) (2016) 51-60.
7. J.-L.K. Kambale, J.A. Asimonyio, R.E. Shutsha, E.W. Katembo, J.M. Tsongo, P.K. Kavira, E.I. Yokana, K.K. Bukasa, H.S. Nshimba, P.T. Mpiana, K.N. Ngbolua. Etudes floristique et structurale des forêts dans le domaine de chasse de Rubi-Télé (Province de Bas-Uélé, République Démocratique du Congo). *International Journal of Innovation and Scientific Research* 24 (2) (2016) 309-321.
8. J.-L.K. Kambale, R.E. Shutsha, E.W. Katembo, J.M. Omatoko, F.B. Kirongozi, O.D. Basa, E.P. Bugenth, E.I. Yokana, K.K. Bukasa, H.S. Nshimba, K.N. Ngbolua. Etude floristique et structurale de deux groupements végétaux mixtes sur terre hydromorphe et ferme de la forêt de Kponyo (Province du Bas-Uélé, R.D. Congo). *International Journal of Innovation and Scientific Research* 24 (2) (2016) 300-308.
9. P.K. Kavira, F.B. Kirongozi, J.-L.K. Kambale, J.M. Tsongo, N.A. Shalufa, K.K. Bukasa, P.Y. Sabongo, H.K. Nzapo, K.N. Ngbolua. Caractéristiques de la régénération naturelle du sous-bois forestier du Jardin botanique S. Lisowski (Kisangani, République Démocratique du Congo). *International Journal of Innovation and Scientific Research* 24 (2) (2016) 322-331.
10. T.B. Mambo, J.U. Thumitho, E.L. Tambwe, C.M. Danadu, J.A. Asimonyio, A.B. Kankonda, J.A. Ulyel, C.M. Falanga, K.N. Ngbolua. Etude qualitative du régime alimentaire de *Hippopotamys psittacus* (Boulenger, 1897: Osteoglossiformes, Mormyridae) du fleuve Congo à Kisangani (RD Congo). *International Journal of Innovation and Scientific Research* 21 (2) (2016) 321-329.
11. F.M. Masudi, A. Dudu, G. Katuala, J.A. Asimonyio, P.K. Museu, B.Z. Gbolo, K.N. Ngbolua. Biodiversité des rongeurs et Soricomorphes de champs de cultures mixtes de la région de Kisangani, République Démocratique du Congo. *International Journal of Innovation and Applied Studies* 14 (2) (2016) 327-339.
12. K.N. Ngbolua, J.A. Asimonyio, N. Ndrodza, B. Mambo, P. Bugenth, Y. Isangi, J.K. Mukirania, L. Ratsina, N.K. Ngombe, P.T. Mpiana. Valeur nutritive et teneur en acide cyanhydrique de huit espèces végétales consommées par *Okapia johnstoni* (Mammalia: Giraffidae) en République Démocratique du Congo. *International Journal of Innovation and Scientific Research* 23 (2) (2016) 419-427.

13. K.N. Ngbolua, B.G. Badjedjea, B.J. Akuboy, M.F. Masudi, J.A. Asimonyio, G.N. Bongo, A.D. Siasia. Contribution to the Knowledge of Amphibians of Kponyo village (DR Congo). *J. of Advanced Botany and Zoology*, V4I1 (2016). DOI: 10.15297/JABZ.V4I1.04.
14. K.N. Ngbolua., A. Mafoto, M. Molongo, G.M. Ngemale, C.A. Masengo, Z.B. Gbolo, P.T. Mpiana, G.N. Bongo. Contribution to the Inventory of "Protected Animals" Sold As Bush Meats in Some Markets of Nord Ubangi Province, Democratic Republic Of The Congo. *J. of Advanced Botany and Zoology*, V3I2 (2015). DOI: 10.15297/JABZ.V3I2.02.
15. K.N. Ngbolua, A. Mafoto, M. Molongo, J.P. Magbukudua, G.M. Ngemale, C.A. Masengo, K. Patrick, H. Yabuda, J. Zama, F. Veke. Evidence of new geographic localization of *Okapia johnstoni* (Giraffidae) in Democratic Republic of the Congo: The rainforest of "Nord Ubangi" district. *Journal of Advanced Botany & Zoology*. V2I1 (2014). DOI: 10.15297/JABZ.V2I1.02.
16. K.N. Ngbolua, G.M. Ngemale., N.F. Konzi, C.A. Masengo, Z.B. Gbolo, B.M. Bangata., T.S. Yangba, N. Gbiangbada. Utilisation de produits forestiers non ligneux à Gbadolite (District du Nord-Ubangi, Province de l'Equateur, R.D. Congo): Cas de *Cola acuminata* (P.Beauv.) Schott & Endl. (Malvaceae) et de *Piper guineense* Schumach. & Thonn. (Piperaceae). *Congo Sciences* 2 (2) (2014) 61-66.
17. M.B. Fenton, N. B. Simmons. *Bats: a world of science and mystery*. The University of Chicago Press: Chicago, USA, 2015.
18. C. Fargeot. Chasse commerciale et le négoce de la venaison en Afrique Centrale forestière. *Game and Wildlife Science* 21 (4) (2004) 817-833.
19. M. Thibault, S. Blaney. The oil industry as an underlying factor in the bushmeat crisis in Central Africa. *Conservation Biology* 17(6) (2003) 1807-1813. DOI 10.1111/j.1523-1739.2003.00159.x.
20. D.A.R. Taylor, B.O. Kankan, M.R. Wagner. The role of the fruit bat, *Eidolon helvum*, in seed dispersal, survival, and germination in *Milicia excelsa*, a threatened West African hardwood. In: Cobbinah, J. R. and Wagner, M. R. (eds): *Research advances in restoration of iroko as a commercial species in West Africa*. Forestry Research Institute of Ghana (FORIG), Kumasi, Ghana: pp. 29-39, 2000.
21. A.O. Kamins, O. Restif, Y. Ntiemoa-Baidu, R. Suu-Ire, D.T.S. Hayman, A.A. Cunningham, J.L.N. Wood, J.M. Rowcliffe. Uncovering the fruit bat bushmeat commodity chain and the true extent of fruit bat hunting in Ghana, West Africa. *Biological Conservation* 144 (12) (2011) 3000-3008. DOI 10.1016/j.biocon.2011.09.003.
22. M.C. Peel, B.L. Finlayson, T.A. McMahon. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11 (2007) 1633-1644. DOI 10.5194/hess-11-1633-2007, 2007.
23. N. Ifuta. Paramètres écologiques et hormonaux durant la croissance et la reproduction d'*Epomops franqueti* (Mammalia: Chiroptera) de la forêt ombrophile équatoriale de Masako (Kisangani-Zaire). PhD thesis, KUL, Leuven: Belgium, 1993.
24. H.H. Nshimba. Etude floristique, écologique et phytosociologique des forêts de l'île Mbiye à Kisangani, RD Congo. Master's Thesis, ULB, Brussels: Belgium, 2008.
25. P.P. Grasse. *Traité de Zoologie, anatomie, systématique, biologie*. Tome XVIII, Fascicule 2. Mammifères. Les ordres: anatomie, éthologie, systématique. Masson et Cie. Paris Vie, 1955.
26. M. Dallimer, T. King, D. Cope, M. Borge Jiana. Estimation of population density of *Eidolon helvum* on the island of Príncipe, Gulf of Guinea. *Mammalia*, 70 (1-2) (2006) 48-51. DOI 10.1515/MAMM.2006.014.
27. D. Thomas, M. Henry. *Eidolon helvum*-African Straw-coloured Fruit Bat. In: Happold, M. and Happold, D.C.D.: *Mammals of Africa. Volume IV: Hedgehogs, Shrews and Bats-Bloomsbury* (2013) 232-234
28. G. Gembu. *Peuplement et dynamique alimentaire des Chiroptères frugivores (Mammalia) de la Réserve Forestière de Yoko (Province Orientale, R. D. Congo)*, Kisangani PhD Thesis. University of Kisangani, 2012.
29. A. Hassanin, S. Khouider, G.C. Gembu, S.M. Goodman, B. Kadjo, N. Nesi, X. Pourrut, E. Nakouné, C. Bonillo. The comparative phylogeography of fruit bats of the tribe Scotonycterini (Chiroptera, Pteropodidae) reveals cryptic species diversity related to African Pleistocene forest refugia. *C.R. Biol.* 338 (2015) 197-211.
30. K. Sabuni. Contribution à l'étude de l'importance relative des mammifères sauvages en général et de *Cephalophus monticola* (Cephalophinés) en particulier dans la consommation de la viande de chasse à Kisangani (Marché du 27 Octobre). Université de Kisangani, Faculté des Sciences, Mémoire, 1978.
31. M. Gambalemoke. Données nouvelles sur la consommation des mammifères sauvages vendus sur le marché central de Kisangani (Haut-Zaïre). Université de Kisangani, Faculté des Sciences, Mémoire, 1989.
32. J.R. Poulsen, C.J. Clark, G. Mavah, P.W. Elkan. Bushmeat supply and consumption in a tropical logging concession in northern Congo. *Conservation Biology*, 23 (6) (2009) 1597-1608. DOI 10.1111/j.1523-1739.2009.01251.x.
33. N.F. Kümpel, E.J. Milner-Gulland, G. Cowlishaw, J.M. Rowcliffe. Incentives for hunting: the role of bushmeat in the household economy in rural Equatorial Guinea. *Human Ecology*, 2 (38) (2010) 251-264. DOI 10.1007/s10745-010-9316-4.
34. A.B. Ononino. *Reglémenter les prélèvements de la viande brousse: entre souhaits et réalités*. Yaoundé, 18 June 2015 ([www.pfbc-cbfp.org/rapports/items/stream2-fr-wildlife.html?file](http://www.pfbc-cbfp.org/rapports/items/stream2-fr-wildlife.html?file), Accessed on 19 February 2017).
35. L. Wetshi, M.N. Biya, J.E. Ruelle. Consommation de mammifères sauvages par la population de Kisangani (Zaire). *Annales de la Faculté des Sciences, Kisangani*, 4 (1987) 135-150.
36. E.J. Milner-Gulland, E.L. Bennett. The SCB 2002 Annual Meeting Wild Meat Group. *Wild meat: the bigger picture*. *Trends in Ecology and Evolution* 18 (7) (2003) 351-357. DOI 10.1016/S0169-5347(03)00123-X
37. IUCN. IUCN red list of threatened species (2010). Version 2010.4. [www.iucnredlist.org](http://www.iucnredlist.org), (accessed on 19.03.11).
38. M.N. Kangoyé, W.Guenda, A. Thiombiano, E.K.V. Kalko, J. Fahr. Les petits mammifères: Chauves-souris / Small mammals: Bats. In: Thiombiano, A. and Kampmann, D. [Eds]: *Biodiversity Atlas of West Africa, Vol. 2: Burkina Faso. Ouagadougou & Frankfurt/Main* (2011) 288-291.
39. R.K.B. Jenkins, A. Keane, A.R. Rakotoarivelo, V. Rakotomboavonjy, F.H. Randrianandrianina, H.J. Razafimanahaka, et al. Analysis of Patterns of Bushmeat Consumption Reveals Extensive Exploitation of Protected

Species in Eastern Madagascar. PLoS ONE 6(12) (2011)  
e27570. doi:10.1371/journal.pone.0027570.

**Citation:** Koto-te-Nyiwa Ngbolua, *et al.* (2017). A survey of the bushmeat trade of the straw-coloured fruit bat (*Eidolon helvum* Kerr, 1792) at Maele Island (Kisangani city, Democratic Republic of the Congo), *J. of Advanced Botany and Zoology*. V4I404. DOI: 10.5281/zenodo.918365.

**Copyright:** © 2017 Koto-te-Nyiwa Ngbolua, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.